

<20% TBSA BURN CHECKLIST

- Burn Consult order placed
- Pictures of burns in chart (can be done by burn team during daytime)
- Perform mapping on Lund-Browder Chart** (see reverse, perform **after** debridement, this will serve as initial and final mapping if discharged)
- If appropriate for discharge, burn-specific discharge instructions and clinic message sent per Burn ED Discharge Guideline
- If requires admission, admit to Burn service as primary team with **Burn A attending** as admitting attending
- If requires admission, place PT/OT consult

>20% TBSA → transport to TBICU before debridement

<20% TBSA → Debride in trauma bay before final disposition

Indications for CT scan:

- Fall >6 feet
- Explosion
- Major traumatic mechanism

- If 7a-3p, coordinate debridement with Burn Dressing Team
- If 3p-12a, trauma chief and TBICU resident will be responsible for debridement
- If 12a-7a, patient will be debrided at ~9AM by Burn Dressing Team

>20% TBSA BURN CHECKLIST

IMMEDIATE TRAUMA BAY INTERVENTIONS

- Start resuscitation sheet
- Record ED arrival time on resuscitation sheet
- Start crystalloid immediately at minimum rate 300 mL/hr
- Insert temperature-sensing foley
- If temperature <37°C, place bair hugger on patient

INITIAL MAPPING AND FLUID CALCULATION (IN TRAUMA BAY, BEFORE DEBRIDEMENT)

- Perform initial mapping using Lund-Browder chart on reverse side
- Calculate initial fluid resuscitation rate (2 mL/kg/%TBSA ÷ 16)
- Record time of initial mapping and calculated fluid resuscitation on resuscitation sheet (Goal = 30 minute from arrival)
- Cover patient with ¾ impervious drape and blankets until initial debridement occurs

TBICU ADMISSION

- Expedite TBICU admission (Goal = 1 hour from arrival)
- Burn service as primary team with Burn A attending as admitting provider
- Record TBICU arrival time on resuscitation sheet
- If >30% TBSA, notify burn attending
- If >30% TBSA, insert Dobhoff tube and initiate tube feeds at 30 kcal/kg/24hr
- Coordinate initial debridement (can be delayed 6-8hr, ensure order for polysporin or bacitracin is placed)
- Give TBICU nurses 2 hours to admit patient before starting debridement. Debridement should not occur within 2 hours of shift change.
- Touch base with ANM prior to beginning debridement to ensure adequate resources
- Place PT/OT Consult

Signature: _____

Date: _____

Draw and shade burned areas of skin (ignore simple erythema)

2°

 3°

Region	%	
	2°	3°
Head		
Neck		
Anterior. trunk		
Posterior. trunk		
Right arm		
Left arm		
Buttocks		
Genitalia		
Right leg		
Left leg		
Total burn		

Signature: _____

Date: _____